NEW TECHNIQUES FOR ANALYZING TIME SERIES AND THEIR IMPORTANCE FOR THE CEOS PROGRAM

First International CEOS Meeting, Sept. 1994

Roy Mendelssohn

PFEG

Monterey, CA

USA

Marie-Hélène Durand

ORSTOM

Paris, FRANCE


UNDERSTANDING CLIMATE CHANGE AND FISHERY DYNAMIC

- Understand How the Environment Affects Fish Dynamics
- Understand What Changes the Climate is Undergoing or is Likely to Undergo
- Combine the Above to Present Scenarios for the Future
- Techniques to be Discussed Mainly Aimed At Second Question


WHAT IS CHANGE?

First International CEOS Meeting, Sept. 1994


Change in the mean level (trend):



Change in the seasonal cycle:



Change in the variance structure:



STATISTICAL CHARACTERIZATION OF CHANGE

First International CEOS Meeting, Sept. 1994

TRENDS

- Global and/or Local variability
- Long-term trend or change in the seasonal component?
- Time and frequency domains can both be considered

NON-STATIONARY SPECTRA

Changes in the variance structure through time

REGIME SHIFTS


- Tests and modeling of shifts between regimes
- Hidden Markov Models


LONG-TERM MEMORY MODELS


- Persistence of events through time
- Related to "chaotic dynamics"


DECOMPOSITION OF A TIME SERIES














EXAMPLE: SST OFF CALIFORNIA COAST




CALCULATION OF TREND











CALCULATION OF SEASONAL







CALCULATION OF SEASONAL (AGAIN)



CALCULATION OF AR









EXAMPLE: N-S WIND STRESS OFF CALIFORNIA





SEASONAL COMPONENT



AR AND ERROR SERIES

25

-25

-75

-125

-175 | 1940

25

-25

-75

-125

-175 –

1940

1950


1960

1970

1980

1990

2000



EXTENSIONS OF METHODLOGY

- Statistical tests of change points, linearity etc.
- Ability to model more exactly abrupt changes
- Multivariate factor model
 - One or more common trends can differ by affine transformation
 - One or more common seasonals can differ by affine transformation
 - Can perform analysis of variance type tests for which model is "best" alternative